



# HHV-6 FOUNDATION

Set:	HHV-7 Infected Cell Lines
Reagent:	SupT-1 / KHR
Virus:	HHV-7 (KHR)
Cell count:	5x10 <sup>6</sup> cells/vial
Contributor:	Yamanishi, K. Laboratory of Virology, Division of Biomedical Research, National Institute of Biomedical Innovation. Japan.
Description:	SupT-1 cells infected with HHV-7 KHR strain. This strain was obtained in 1994 from a pediatric patient with exanthen subitum by Tanaka et al.
References:	<p>Tanaka K, Kondo T, Torigoe S, Okada S, Mukai T, Yamanishi K. Human herpesvirus-7: another causal agent for roseola (exanthen subitum). J Pediatr. 1994 Jul;125(1):1-5.</p> <p>Smith SD, Shatksy M, Cohen PS, Warnke R, Link MP, Glader BE. Monoclonal antibody and enzymatic profiles of human malignant T-lymphoid cells and derived cell lines. Cancer Res. 1984 Dec;44(12 Pt 1):5657-60.</p>
Special Notes:	The SupT-1 cell line was originally derived in 1984 from a pediatric patient with T cell lymphoblastic lymphoma by Smith et al.
NIH Status:	Not acquired
MTA Required:	No